REFERENCES

Aimes, H., Bokonon-Gantaa, Grooteb,H., and Neuenschwander,P.(2002). Socio-economic impact of biological control of mango mealy bug in Benin, Ecosystems and Environment, Vol. 93:367–378.

Beardsley, J., German, W., Reimer, T., and Sanford, R. (1998). Mealy bug wilt, mealy bugs, and ants of pineapple, Plant Disease, Vol. 72 No.7:558-565.

Buss, E., and Turner, Jay. (2006). Scale insect and mealy bugs on ornamental plants, Institute of Food and Agriculture Sciences, ENY-323.

Blumberg, D., and Drieschet, R., (2001). Encapsulation Rates of three encyrtid parasitoids by three mealy bug species (Homoptera: Pseudococcidae) Found commonly as pests in commercial greenhouses biological control Vol.22: 191–199.

Clark, J., and Furnees, G.(1992). Common green house pest-mealy bugs, The Biology of Grass house Pest and Their Nature Enemies.

Ceballo, F.A. & Walter, G.H. (2004). Life history parameters and bio control potential of the mealy bug parasitoid *Coccidoxenoides peregrinus* (Timberlake) (Hymenoptera: Encyrtidae): asexuality, fecundity and ovipositional patterns. Biological Control Vol.29: 235–244.

Charles, J. (2008). A survey of mealy bugs and their natural enemies in horticultural crops in north island, implications for biological control**:** 405-418.

Cudjoe, A., Neuenschwander, P., And Copland, M.(1993). Interference by ants in biological control of the cassava mealy bug *Phenacoccus manihoti* (Hemiptera: Pseudococcidae) in Ghana, Bulletin of Entomological Research, Volume No. 83(01):15-22.

Daane, K., Almeida, R., Bell, V., Walkar, J., Bottom, M., Fallahzadeh, M., Mani, M., Miano, J., Forza, S., Walton, V., and Zaviero, T. (2012). Biology and management of mealy bugs in vineyards arthropods management in vineyards, 271-307

Davies, A.P., Ceballo, F.A. & Walter, G.H.(2004). Is the potential of *Coccidoxenoides perminutus*, a mealy bug parasitoid, limited by climatic or nutritional factors? Biological Control Vol.31: 181–188.

Goble, H. Wukasch , R . , and Sabourin , M . (2012). Scale insect and mealy bugs on house plants, Pest diagnostic Clinic.

Groote, H., Neuenschwander, P., and Bokonon-Ganta, A. (2002). Socio-Economic Impact of Biological Control of Mango Mealy bug In Benin, Vol.93**:** 367-378.

Godfray, H., and Waage, J. (1991). Predictive Modelling In Biological Control: The Mango Mealy Bug (*Rastrococcus Invadens*) And Its Parasitoids. Journal of Applied Ecology, Vol. 28 (2): 434-453.

Gutierrez, A., Daane, K., Ponti, L., Walton, V., and Ellis, C. (2007). Prospective evaluation of the biological control of vine mealy bug: Refuge Effects and Climate Article.Vol.45 (2): 524-536.

Geiger, C., and Daane, K. (2001). Seasonal movement and sampling of the grape mealy bug, *Planococcus maritimis* (Ehrhorn) (Homoptera: Pseudococcidae) in San Joaquin Valley vineyards. Journal of Economic Entomology Vol.94: 291–301.

Hollingsworth, R. (2005). Limonene ,a citrus extract,for control of mealy bug and scale insect, J. Econ, Entomology. Vol.98 (3) : 772-779.

Johnson, P. (2010). Mealy bugs on Orchids, Insect Research Collection. Orchid digest, Vol.74 (3):170-177.

Lohr, B., Varela, A., and Santos, B. (1990). Exploration for Natural Enemies Of The Cassava Mealy bug, Phenacoccus Manihoti (Homoptera:Pseudococcidae), Biological Control Of This Introduced Pest In Africa Vol.80 (4): 417-425.

Michaud, J. (2012). Three target of classical biological control in the Caribbean : Success, contribution, and failure, International symposium on Biological control of arthropods.

Muniappan, R., Meyerdirk, D., Sengebau, F., Berringer, D., and Reddy, G.(2006). Classical Biological Control of the Papaya Mealy bug, *Paracoccus Marginatus* (Hemiptera: Pseudococcidae) In the Republic of Palau, 89 (2): 212-217.

Neuenschwander, P., Boavida, C., Bokonon-Ganta, A., Gado, A., and Herren, H.(1994). Establishment And Spread of Gyranusoidea Tebyginoyes And Anagyrus Mangicola Noyes (Hymenoptera: Encyrtidae), Two Biological Control Agents Released Against mealy bugs,Vol.No 4(1): 61-69.

Neuenschwander, P., Borowka, R., Phiri, G., Hammans, H., Nyirenda, S., Kapeya, E., and Gadabu, A. (1990). Biological Control Of The Cassava Mealy bug phenacoccus Manihoti (Hom. Pseudococcidae) Byepidinocarsis Lopezi (Hym., Encyrtidae) In Malawi Vol.35(4) :515-526.

Reimer, N., and Johnson, M. (1999). Survey of the natural enemies of Dysmicoccus mealy bugs on pineapple in Hawaii, Volume 44, (1): 47-58.

Shahina, F. , Kazmi , A., Tabassum, , Salma J. And Mahreen, G., (2011) Pathogenecity Of Xenorhabdus Nematophila (Strain Ham-10) Against Wax Moth Larvae And Vine Mealy Bug In Pakistan ,National Nematological Research Centre, Vol.29 (1) :15-23.

Singh,A., Kataria,R., and Kumar,D. (2012). Repellence property of traditional plant leaf extracts against *Aphis gossypii* Glover and *Phenacoccus solenopsis* Tinsley, African Journal of Agricultural Research, Vol. 7(11):1623-1628.

Tansuri, B. (2010). Adoption of biochemical agents at the field level for management of mealy bugs challenges and proposed solution, Journal of bio pesticides.Vol.3 (1): 055-057.

Walton, V., and Pringle, K. (1999). Effects of pesticides used on table grapes on the mealy bug parasitoid *Coccidoxenoides perigrinus* (Timberlake) (Hymenoptera: Encyrtidae). South African Journal of Enology and Viticulture Vol.20:31–34.

Watson, G.W., and Kubirita, J. (2005). Identification of mealy bugs (Hemiptera: Pseudococcidae) on banana and plantain in Africa. African Entomology Vol.13: 35–47.

Zeddies, J., Schaab, R., Neuenschwander, P., Herren, H. (2005). Economics of biological control of cassava mealy bug in Africa.